

Fixed Point Theorems in Parametric Metric Space for Contraction Mappings

Ria Sharma

Department of Mathematics, Chandigarh University, Punjab, India

ABSTRACT

Parametric Metric Space (P.M.S) is a well known generalization of Metric Space (MS) which was introduced and studied by Hussian (a new approach to M.S)in 2014. In the present paper, we extended fixed point theorems (F.P.T)of Zamfirescu and Hardy –Rogers based on injective mapping to Parametric Metric Space (P.M.S) using contraction conditions. Moreover, we represented an examples to validate our result.

Keyword: Fixed point; Contraction mapping; Metric Space; Complete Parametric Metric Space; injective mapping

I. INTRODUCTION

Real analysis is the most important branch of mathematics. Among several branches of real analysis, functional analysis is the most important part of real analysis. Functional analysis is divided into two parts: linear and non-linear. F.P.T is an important part of non-linear functional analysis since 1960. F.P.T has various applications in field of pure and applied mathematics as well as in physical, economic, differential equation, integral equation and life sciences. It has emerged as one of the major links between abstract mathematics and its applications. It is used in differential equations, integral theory, artificial intelligence, computer science, decision making, medical diagnosis, neural network, social science and many other related areas. F.P.T deals with the classical approach to the exact solution and to check the stability of the system. F.P.T has fascinated lots of researchers since 1922, with the celebration of Banach (Polish mathematician) contraction principle [25] which provided a constructive method to find a fixed point of a mapping. Banach contraction principle states: A mapping $T: X \rightarrow X$ defined on M.S

(X, δ) is called contraction if $\delta(Ta, Tb) \leq k\delta(a, b) \forall a, b \in X \text{ \& } 0 < k < 1$. Since then this theorem became an important tool for the development of nonlinear analysis. Contraction conditions begins by studying Banach contraction principle which are utilized in various F.P.T for some generalized M.S. Various F.P.T were attained by expanding Banach contraction principal [2,3,4,17,18, 21, 23, 24, 25].

Kannan [16] proved that a mapping $T: X \rightarrow X$ defined on M.S (X, δ) is called contraction mapping if $\delta(Ta, Tb) \leq k[\delta(a, Ta) + \delta(b, Tb)] \forall a, b \in X \text{ \& } 0 < k < 1/2$ for operators that need not be continuous. Further, Chatterjea [4], proved a F.P.T for discontinuous mapping, which is actually a kind of dual of Kannan mapping. Some of the generalizations of M.S are cone metric spaces, partial metric spaces, P.M.S, “parametric b-metric space” (P.B.M.S) etc. [1,2,7,11,12,14,15,17,19,22,23,29,30] introducing and modifying the metric axioms.

Hussain et al. [13] studied and introduced the notion of P.M.S, and later on gave generalized P.M.S and introduced P.B.M.S which is combination of M.S as well as b-M.S [14]. In the present paper we proved fixed point theorems on P.M.S for expansive mapping. We generalized the fixed point theorems of Zamfirescu [36] and Hardy–Rogers [13] to their parametric versions in complete metric space.

Definition 1.1: Let X be a non empty set and $T_p: X \times X \times (0, \infty) \rightarrow (0, \infty)$ be a map on X such that $\forall a, b, c \in X$ and $t > 0$

- (a). $T_p(a, b, t) = 0$ if and only if $a = b$
- (b). $T_p(a, b, t) = T_p(b, a, t)$
- (c). $T_p(a, b, t) \leq T_p(a, c, t) + T_p(c, b, t)$

Then pair (X, δ) is called P.M.S.

Definition 1.2: Let X be a non empty set and (X, δ) be a P.M.S and let $\{a_i\}$ be a sequence in X

- (a) If $\log_{i \rightarrow \infty}(a_i, a, t) = 0 \Rightarrow \log_{i \rightarrow \infty} a_i = a$, for all $t > 0$ then sequence $\{a_i\}_{i=1}^{\infty}$ converges $a \in X$
- (b) If $\log_{n \rightarrow \infty}(a_i, a_j, t) = 0$ for all $t > 0$ then sequence $\{a_i\}_{i=1}^{\infty}$ is called Cauchy sequence.
- (c) If every Cauchy sequence is convergent, then P.M.S (X, δ) is a complete P.M.S.

(d) Let (X, δ) be a P.M.S and $T_p: X \rightarrow X$ be a mapping, then We say T is a continuous mapping in X , if for any sequence $\{a_i\}_{i=1}^{\infty} \in X$ such that $\log_{i \rightarrow \infty} a_i = x \Rightarrow \log_{i \rightarrow \infty} T_p a_i = T_p a$

II. MAIN RESULT:

Theorem 2.1: Let (X, δ) be a complete P.M.S and $T_p: X \rightarrow X$ be an injective mapping satisfying the condition (2.1.1)

$$\delta(T_p a, T_p b, t) \leq \alpha_1 \delta(a, b, t) + \alpha_2 \delta(a, T_p a, t) + \alpha_3 \delta(b, T_p b, t) + \alpha_4 \delta(a, T_p b, t) + \alpha_5 \delta(b, T_p a, t)$$

$\forall t \in [0, 1); \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5 > 0; a, b \in X$ & $a \neq b$ have a fixed point if $\alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 < 1$ and moreover a unique point if $\alpha_1 + \alpha_4 + \alpha_5 < 1$.

Proof: Let $a_0 \in X$, Define iterative sequence $\{a_i\}_{i=1}^{\infty}$ follows: $T_p a_i = a_{i+1}$ for $i = 1, 2, 3, \dots$. If for some i ,

$T_p a_i = a_i$, then a_i is the fixed point. Otherwise $T_p a_i \neq a_i$, using inequality (2.1.1)

$$\begin{aligned} \delta(a_{i+1}, a_{i+2}, t) &= \delta(T_p a_i, T_p a_{i+1}, t) \\ &\leq \alpha_1 \delta(a_i, a_{i+1}, t) + \alpha_2 \delta(a_i, T_p a_i, t) + \alpha_3 \delta(a_{i+1}, T_p a_{i+1}, t) + \alpha_4 \delta(a_i, T_p a_{i+1}, t) \\ &\quad + \alpha_5 \delta(a_{i+1}, T_p a_i, t) \end{aligned}$$

$$\leq \alpha_1 \delta(a_i, a_{i+1}, t) + \alpha_2 \delta(a_i, a_{i+1}, t) + \alpha_3 \delta(a_{i+1}, a_{i+2}, t) + \alpha_4 \delta(a_i, a_{i+2}, t) + \alpha_5 \delta(a_{i+1}, a_{i+1}, t)$$

$$\leq \alpha_1 \delta(a_i, a_{i+1}, t) + \alpha_2 \delta(a_i, a_{i+1}, t) + \alpha_3 \delta(a_{i+1}, a_{i+2}, t) + \alpha_4 [\delta(a_i, a_{i+1}, t) + \delta(a_{i+1}, a_{i+2}, t)]$$

$$(1 - \alpha_3 - \alpha_4) \delta(a_{i+1}, a_{i+2}, t) \leq (\alpha_1 + \alpha_2 + \alpha_4) \delta(a_i, a_{i+1}, t)$$

$$\delta(a_{i+1}, a_{i+2}, t) \leq \frac{(\alpha_1 + \alpha_2 + \alpha_4)}{(1 - \alpha_3 - \alpha_4)} \delta(a_i, a_{i+1}, t)$$

$$\delta(a_{i+1}, a_{i+2}, t) \leq k \delta(a_i, a_{i+1}, t)$$

$$\text{Where } k = \frac{(\alpha_1 + \alpha_2 + \alpha_4)}{(1 - \alpha_3 - \alpha_4)} < 1 \Rightarrow \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 < 1$$

Continuing iterations up to i times

$$\delta(a_{i+1}, a_{i+2}, t) \leq k^i \delta(a_0, a_1, t)$$

As we know if $\{a_i\}_{i \rightarrow \infty}$ be a sequence in P.M.S (X, δ) such that $\delta(a_{i+1}, a_{i+2}, t) \leq k^i \delta(a_0, a_1, t)$

$\forall t \in [0,1] \& i = 1,2,3, \dots$ then $\{a_i\}_{i \rightarrow \infty}$ is a Cauchy sequence in (X, δ) . Since (X, δ) is a complete P.M.S; $\{a_i\}_{i \rightarrow \infty}$ converges. Let $a^* \in X$, then $\lim_{n \rightarrow \infty} x_i \rightarrow x^*$. Again T_p is continuous, therefore

$$T_p a^* = T_p(\lim_{n \rightarrow \infty} a_n) = \lim_{n \rightarrow \infty} T_p a_i = a^* \Rightarrow T_p a^* = a^*$$

Implies T_p has a fixed point $T_p a^* = a^*$ in X .

Now we will show that a^* is unique. for that suppose b^* is another fixed point therefore $T_p b^* = b^*$. Therefore by inequality (2.1.1) we have

$$\delta(T_p a^*, T_p b^*, t) \leq \alpha_1 \delta(a^*, b^*, t) + \alpha_2 \delta(a^* T_p a^*, t) + \alpha_3 \delta(b^*, T_p b^*, t) + \alpha_4 \delta(a^*, T_p b^*, t) + \alpha_5 \delta(b^*, T_p a^*, t)$$

$$\delta(a^*, b^*, t) \leq \alpha_1 \delta(a^*, b^*, t) + \alpha_2 \delta(a^*, b^*, t) + \alpha_3 \delta(b^*, b^*, t) + \alpha_4 \delta(a^*, b^*, t) + \alpha_5 \delta(b^*, a^*, t)$$

$$\delta(a^*, b^*, t) \leq \alpha_1 \delta(a^*, b^*, t) + \alpha_4 \delta(a^*, b^*, t) + \alpha_5 \delta(a^*, b^*, t)$$

$$\delta(a^*, b^*, t) \leq (\alpha_1 + \alpha_4 + \alpha_5) \delta(a^*, b^*, t)$$

$$(1 - \alpha_1 - \alpha_4 - \alpha_5) \delta(a^*, b^*, t) \leq 0$$

$\Rightarrow a^* = b^*$. Since $\alpha_1 + \alpha_4 + \alpha_5 < 1$ implies that a^* and b^* are not different point but are same.

Hence a^* is unique.

Theorem 2.2: Let (X, T_p) be a complete P.M.S and $T_p: X \rightarrow X$ be an injective mappings satisfying condition

$$(2.2.1) \quad (T_p a, T_p b, t) \leq \alpha \text{ Max} \left\{ \delta(a, b, t), \frac{\delta(a, T_p a, t) + \delta(b, T_p b, t)}{2}, \frac{\delta(a, T_p b, t) + \delta(b, T_p a, t)}{2} \right\}$$

$\forall t \in [0,1]; \alpha, \beta, \gamma > 0; x, y \in X \& a \neq b$ then T_p has a fixed point if $\alpha + \beta + \gamma < 1$ and moreover a unique fixed point if $\alpha + \gamma < 1$.

Proof: Let $a_0 \in X$, Define iterative sequence $\{a_i\}_{i=1}^{\infty}$ follows: $T_p a_i = a_{i+1}$ for $i = 1, 2, 3, \dots$. If for some i ,

$T_p a_i = a_i$, then a_i is the fixed point. Otherwise $T_p a_i \neq a_i$, using inequality (6.1)

$$\delta(a_{i+1}, a_{i+2}, t) = \delta(T_p a_i, T_p a_{i+1}, t)$$

$$\begin{aligned} &\leq \alpha \text{Max}\left\{\delta(a_i, a_{i+1}, t), \frac{\delta(a_i, T_p a_i, t) + \delta(a_{i+1}, T_p a_{i+1}, t)}{2}, \frac{\delta(a_i, T_p a_{i+1}, t) + \delta(a_{i+1}, T_p a_i, t)}{2}\right\} \\ &\leq \alpha \text{Max}\left\{\delta(a_i, a_{i+1}, t), \frac{\delta(a_i, a_{i+1}, t) + \delta(a_{i+1}, a_{i+2}, t)}{2}, \frac{\delta(a_i, a_{i+2}, t) + \delta(a_{i+1}, a_{i+1}, t)}{2}\right\} \\ &\leq \alpha \text{Max}\left\{\delta(a_i, a_{i+1}, t), \frac{\delta(a_i, a_{i+1}, t) + \delta(a_{i+1}, a_{i+2}, t)}{2}, \frac{\delta(a_i, a_{i+1}, t) + \delta(a_{i+1}, a_{i+2}, t)}{2}\right\} \\ &\Rightarrow \delta(a_{i+1}, a_{i+2}, t) \leq \alpha \text{Max}d(a_i, a_{i+1}, t) \end{aligned}$$

Therefore, by successive iteration

$$\delta(a_{i+1}, a_{i+2}, t) \leq \alpha^i \delta(a_0, a_1, t)$$

As we know if $\{a_i\}_{i \rightarrow \infty}$ be a sequence in P.M.S (X, δ) such that $\delta(a_{i+1}, a_{i+2}, t) \leq \alpha^i \delta(a_0, a_1, t)$

$\forall t \in [0,1] \& i = 1,2,3, \dots$ then $\{a_i\}_{i \rightarrow \infty}$ is a Cauchy sequence in (X, δ) . Since (X, δ) is a complete P.M.S; $\{a_i\}_{i \rightarrow \infty}$ converges. Let $a^* \in X$, then $a \rightarrow a^*$. Again T_p is continuous, therefore

$$T_p a^* = T_p (\lim_{n \rightarrow \infty} a_i) = \lim_{n \rightarrow \infty} T_p a_i = a^* \Rightarrow T_p a^* = a^*$$

Implies T_p has a fixed point $T_p a^* = a^*$ in X.

Now we will show that a^* is unique. For that suppose y^* is another fixed point therefore $T_p y^* = y^*$. Therefore by inequality (2.2.1) we have

$$\begin{aligned} &\delta(T_p a^*, T_p b^*, t) \\ &\leq \alpha \text{Max}\left\{\delta(a^*, b^*, t), \frac{\delta(a^*, T_p a^*, t) + \delta(b^*, T_p b^*, t)}{2}, \frac{\delta(a^*, T_p b^*, t) + \delta(b^*, T_p a^*, t)}{2}\right\} \\ &\delta(T_p a^*, T_p b^*, t) \leq \alpha \text{Max}\left\{\delta(a^*, b^*, t), \frac{\delta(a^*, a^*, t) + \delta(b^*, b^*, t)}{2}, \frac{\delta(a^*, b^*, t) + \delta(b^*, a^*, t)}{2}\right\} \end{aligned}$$

$$\delta(T_p a^*, T_p b^*, t) \leq \alpha \text{Max}\{\delta(a^*, b^*, t)\}$$

$$\delta(a^*, b^*, t) \leq \alpha \delta(a^*, b^*, t)$$

$$\Rightarrow (1 - \alpha) \delta(a^*, b^*, t) \leq 0$$

$$\Rightarrow \delta(a^*, b^*, t) = 0 \text{ since } \alpha > 1 \Rightarrow a^* = b^*$$

Hence T_p has a unique point.

Example 2.3: Let (X, δ) be a complete P.M.S, where $T_p: R^+ \rightarrow R^+$ is a mapping defined as $\delta(a, b, t) = t|a - b|^q$ such that $a_n = 1 + \frac{1}{n}$ and $b_n = 1 + \frac{2}{n}$, therefore

$$\begin{aligned}\delta(a_n, b_n, t) &= t|a_n - b_n|^q = t \left| \left(1 + \frac{1}{n}\right) - \left(1 + \frac{2}{n}\right) \right|^q = t \left| \frac{1}{n} - \frac{2}{n} \right|^q = t \left| -\frac{1}{n} \right|^q = t \left(\frac{1}{n} \right)^q \\ &= t \frac{1}{n^q}\end{aligned}$$

$$\log_{n \rightarrow \infty} \delta(a_n, b_n, t) = \log_{n \rightarrow \infty} t \frac{1}{n^q} = t \log_{n \rightarrow \infty} \frac{1}{n^q} = 0 \text{ for } t > 0$$

$$\Rightarrow \log_{n \rightarrow \infty} \delta(a_n, b_n, t) \rightarrow 0$$

as both $a_n = 1 + \frac{1}{n}$ and $b_n = 1 + \frac{2}{n}$ tends to 1 as $n \rightarrow \infty$. Hence 1 is the fixed point.

It satisfies all the conditions of complete parametric metric space for $t > 0$ and theorems [2.1,2.2]

III. CONCLUSION:

The aim of this paper is to verify that existing theorems in complete metric space are true or not for parametric metric space if a parameter $t > 0$ is added under the contraction conditions.

IV. REFERENCES :

1. Aage and Salunke: Some fixed point theorems for expansion onto mappings on cone metric spaces, Acta Math. Sin. Engl. Ser., 27(6)(2011), 1101-1106
2. Alghamdi, Hussain and Salimi: Fixed point and coupled fixed point theorems on b-metric-like spaces, J. Inequal. Appl., 2013(2013)
3. Bari and Vetro: Fixed points, attractors and weak fuzzy contractive mappings in a fuzzy metric space, Journal of Fuzzy Mathematics, 13(4)(2005), 973982.
4. Chatterjea, S. K., Fixed-point theorems, C. R. Acad. Bulgare Sci., vol. 25, pp. 727-730, 1972.
5. Czerwik.S , Contraction mappings in b-metric spaces , Acta Math. Inf. Univ. Ostravensis, 1(1993),5-11.
6. Czerwik :Nonlinear set-valued contraction mappings in b-metric spaces, Atti Sem. Mat. Fis. Univ. Modena, 46(1998), 263-276.

7. Daheriya, Shrivastava and Ughade: Parametric Metric Space, Parametric b-metric Space and Expansive Type Mapping ,International Journal of Mathematics And its Applications,Volume-4 (2016),107-117
8. Daheriya, Jain and Ughade: Some Fixed Point Theorem for Expansive Type Mapping in Dislocated Metric Space, ISRN Mathematical Analysis, 2012(2012), Article ID 376832.
9. Daffer and Kaneko: On expansive mappings, Math. Japonica., 37(1992), 733-735.
10. George and Veeramani.: On some results in fuzzy metric spaces, Fuzzy Sets and Systems, 64(3)(1994), 395399.
11. Grabiec: Fixed points in fuzzy metric spaces, Fuzzy Sets and Systems, 27(3)(1988), 385389
12. Huang, Zhu and Wen :Fixed point theorems for expanding mappings in partial metric spaces, An. St. Univ. Ovidius Constant a, 20(1)(2012), 213-224.
13. Hussain,Khaleghizadeh, Salimi and Abdou: A New Approach to Fixed Point Results in Triangular Intuitionistic Fuzzy Metric Spaces, Abstract and Applied Analysis, 2014(2014), Article ID 690139.
14. Hussain Nawab and SalimibPeyman and ParvanehcVahid; Fixed point results for various contractions in parametric and fuzzy b-metric spaces , J. Nonlinear Sci. Appl. 8 (2015), 719–739
15. Kannan R., Some results on fixed points, Bull. Calcutta Math. Soc., vol. 60, pp.71-76, 1968.
16. Kramosil and Michalek:Fuzzy metrics and statistical metric spaces, Kybernetika, 11(5)(1975), 336-344.
17. Shatanawi and Awawdeh: Some fixed and coincidence point theorems for expansive maps in cone metric spaces, Fixed Point Theory and Applications, 2012(2012).
18. Jain,Daheriya and Ughade: Fixed Point, Coincidence Point and Common Fixed Point Theorems under Various Expansive Conditions in Parametric Metric Spaces and Parametric b-Metric Spaces, Gazi University Journal of Science, 2016 , 95-107 ,volume 9
19. Jain, Daheriya and Ughade: Fixed Point, Coincidence Point and Common Fixed Point Theorems under Various Expansive Conditions in b-Metric Spaces, International Journal of Scientific and Innovative Mathematical Research, 3(9)(2015), 26-34.
20. Mishra.S.N,N.SharmaHardy.GE, Rogers. TD: A generalization of a fixed point theorem , Reich. Can. Math. Bull. **16**, 201-206 (1973)
21. andS.L.Singh,Common fixed point of mappings on fuzzy metric spaces, Internat.J.Math.Math.Sci, 17(1994),253-258.
22. Muraliraj and Hussain :Coincidence and Fixed Point Theorems for Expansive Maps in d-Metric Spaces, Int.Journal of Math. Analysis, 7(44)(2013), 21712179.
23. Mustafa, Awawdeh and Shatanawi: Fixed Point Theorem for Expansive Mappings in G-Metric Spaces, Int. J.Contemp. Math. Sciences, 5(50)(2010), 2463-2472.
24. Pachpatte.B.G, On certain fixed point mappings in metric spaces, Journal of the Maulana Azad College of Technology, vol. 13, pp. 59–63, 1980.
25. Palais.Richard S, A simple proof of the banach contraction principle,J. fixed point theory appl. 2 (2007), 221–223.
26. Park ,Intuitionistic fuzzy metric spaces, Chaos, Solitons and Fractals, 22(5)(2004), 10391046.
27. Pant, some fixed point theorems in fuzzy metric spaces, TamkangJ.Math. 1(2009),59-66

28. Rus.L.A, Metric space with fixed point property with respect to contractions, *Studia Univ. Babeş-Bolyai*, 51(2006), 115-121.
29. Saluja,Dhakda and Magarde, Some fixed point theorems for expansive type mapping in dislocated metric space, *Mathematical Theorey and Modeling*, 3(2013)
30. Shatanawi and Awawdeh,Some fixed and coincidence point theorems for expansive maps in cone metric spaces, *Fixed Point Theory and Applications*, 2012(2012).
31. Vasuki, A common fixed point theorem in a fuzzy metric space,*Fuzzy sets and systems*, 97(1998),395-397.
32. Vetro.Gopal and M.Imdad, Common fixed point theorems (*J.Math*,52(3),(2010),573-590.
33. Wang, Li, Gao and Iseki: Some fixed point theorems for expansion mappings, *Math. Japonica.*, 29(1984), 631-636.
34. Wang, S.Z., Li, B.Y., Gao, Z.M., Iseki, K., Some fixed point theorems for expansion mappings, *Math. Jaon* 29(1984), 631- 636.
35. Zadeh: Fuzzy sets, *Information and Computation*, 8(1965), 338353
36. Zamfirescu.T, Fixed point theorems in metric spaces, *Arch. Math. (Basel)*, 23 (1972), pp. 292-298