

Emphasis of Coefficients on the Convergence Rate of Fixed Point Iterative Algorithm in Banach Space

Naveen Kumar¹, Surjeet Singh Chauhan (Gonder)²

Department of Mathematics

University Institute of Sciences (UIS), Chandigarh

University, Gharuan, Mohali-140413, Punjab, India

ABSTRACT

This paper deals with the fixed point iterative schemes that are being utilized to solve the systems of nonlinear equations in various fields and spaces. The convergence speed of iterative processes is highly focused, calculated and compared with various original methods to check the efficiency of these methods towards its optimal solution. The effect on the speed of convergence by the coefficients included in these iterative algorithms are investigated and analyzed in this manuscript. Moreover, the results on the convergence rate of various fixed point iterative schemes are supported by comparing the convergence rate of these iterative plans using the methodology of interchanging the coefficients of these algorithms. The convergence behaviour of these iterative processes is also shown graphically. In a nut shell, the comparison analysis shows that the coefficients involved in such type of schemes may vary the convergence rate of the schemes towards their fixed points.

Keyword: Banach Space, Contractive Mapping, Convergence Rate, Fixed Point Iterations, Fixed Point.

I. INTRODUCTION

This paper deals with the fixed point iteration methods that utilized to solve linear and nonlinear systems. It is assumed that each problem of can be expressed using the fixed point algorithm. Fixed Point Algorithm (FPA) is a type of tools which is available everywhere and use to find more accurate solutions to non-linear systems used in Analysis, Algebra, Geometry, Logic etc. The convergence speed of these procedures play an effective role in the solution of non-linear systems. This paper discusses about the fixed point iterative schemes with it various applications in every field science and technology. A very common and fundamental application

of fixed point in computer science is in recursion theory and mathematical model loops. The utilize of its algorithm is because of the semantics of recursion that are depicted by fixed points of functions. One of the first occurrence of fixed points is in the field of automata (robotics). The theorem about the existence & axioms of the FPs is popularly regarded as the 'fixed point theorem' (FPT). Several remarks, results and consequences in the theory of automata are obtained from the basic results of fixed points.

Several prominent applications related to the fixed point theory (FPT) are there in every field of Mathematical science and technology. In analysis, fixed point has aim perative job in software engineering and in data-flow analysis as programming languages compilers use fixed point calculation for program statement analysis which is often required for code optimization. In numerical analysis, one can calculate the FPs of the iteration functions using FP-iteration. The fixed point iteration for a point τ in the domain of function φ is defined as $\tau_{p+1} = \varphi(\tau_p)$ for $p \geq 0$. In Numerical analysis, it is a method of determining fixed point by doing a number of iteration steps to the function. The fixed point iteration process may be called as a chain point, because the current result of the immediate solution treated as an input to next solution. So it creates a chain point in more appropriate values of the solution.

The Fixed Point Algorithm (FPA) generates a recursive sequence that find the fixed point for some standard functions. The major advantages and benefits of these algorithms are that these are not difficult to implement on Mathematical models and functions. The Fixed Point Algorithm (FPA) uses a value, ideally chosen very close to the fixed point that we want to find and a function $\varphi(\tau)$ that generates a recursively defined sequence τ_p for $p = 0$ and $\tau_{p+1} = \varphi(\tau_p)$ where $p \geq 0$.

The Carl Gustav Jacob's (CGJ) or Jacobi iterative procedure (JIP) is a prominent method for finding solution of diagonal system of equations. Similarly, Gauss Seidel Iteration method (GSIM) utilizes the latest approximate values and scan the mesh points symmetrically from left rows to right rows along with successive rows.

1. Preliminaries

A fixed point of a function $\varphi(\tau)$ is a value τ_0 in the domain of function such that $\varphi(\tau) = \tau$. Geometrically, a fixed point occurs at which the graph of the function $\varphi = \varphi(\tau)$ crosses the graph of $\rho = \tau$.




Fig. 1: Fixed Point

Following are some basic definitions and preliminaries that are used in this paper.

Definition 2.1 Let \square be a non-void set and φ be a self-mapping on \square . A FP of φ is an object in \square that is drawn to itself, i.e. $\tau \in \square$ implies $\varphi(\tau) = \tau$.

Definition 2.2 Assume that $C' \subseteq \square$ be a non-void closed and convex subset. A function $\varphi: C' \rightarrow C'$ is non-expansive if $\|\varphi\tau - \varphi\rho\| \leq \|\tau - \rho\| \forall \tau, \rho \in C'$.
(1)

Definition 2.3. Let (\square, ω) and $\varphi: \square \rightarrow \square$ be a map on \square . Define $F\varphi = \{\tau \in \square \mid \varphi\tau = \tau\}$ as the collection of fixed points of φ , then $\omega(\varphi\tau, \varphi\rho) \leq \mu \omega(\tau, \rho) \forall \tau, \rho \in \square$ and $\mu \in [0, 1)$ is known as Banach contraction's condition. (2)

Definition 2.4. A mapping $\varphi: \square \rightarrow \square$ is supposed to be nonexpansive if $\omega(\varphi\tau, \varphi\rho) \leq \omega(\tau, \rho) \forall \tau, \rho \in \square$. It is called a contraction if for all $\tau, \rho \in \square$, $\exists \mu \in (0, 1)$ such that

$$\|\varphi\tau - \varphi\rho\| \leq \mu \|\tau - \rho\| \tag{3}$$

Definition 2.5. Let $\{\sigma_p\}$ and $\{\sigma'_p\}$ be two sequences approaches to n and m , then $\{\sigma_p\}$ moves faster than $\{\sigma'_p\}$ if $\lim_{p \rightarrow \infty} \frac{\|\sigma_p - n\|}{\|\sigma'_p - m\|} = 0$. (4)

In the following section, we discuss the comparison of various possible iterations obtained by exchanging the coefficients and their comparison analysis is recorded as improved version.

2. Numerical Convergence

Example 3.1: Let $\square = [1, 50]$ with initial values $\sigma_0 = 20$ and $\tau_0 = 30$ where the coefficient $\theta_p = 0.70$ and $\delta_p = 0.80$ for all $p \geq 0$. Define two maps $\varphi_1: \square \rightarrow \square$ and $\varphi_2: \square \rightarrow \square$ by $\varphi_1(\sigma) = \sigma^{\frac{1}{2}}$ and $\varphi_2(\sigma) = \sigma^{\frac{1}{3}}$

where φ_1 and φ_2 are the contraction maps.

Sol. Consider the following fixed point iteration scheme:

$$\begin{aligned} \varphi_2 \sigma_{p+1} &= (1 - \theta_p) \varphi_2 \sigma_p + \theta_p \varphi_1 \tau_p \\ \varphi_2 \tau_p &= (1 - \delta_p) \varphi_2 \sigma_p + \delta_p \varphi_1 \sigma_p \end{aligned} \tag{5}$$

where $\sigma_0 = 20$, $\tau_0 = 30$, $\theta_n = 0.70$ and $\delta_0 = 0.80$ for all $p \geq 0$.

$$\begin{aligned} \varphi_2 \sigma_{p+1} &= \theta_p \varphi_2 \sigma_p + (1 - \theta_p) \varphi_1 \tau_p \\ \varphi_2 \tau_p &= \delta_p \varphi_2 \sigma_p + (1 - \delta_p) \varphi_1 \sigma_p \end{aligned} \tag{6}$$

$$\begin{aligned} \varphi_2 \sigma_{p+1} &= \theta_p \varphi_2 \sigma_p + (1 - \theta_p) \varphi_1 \tau_p \\ \varphi_2 \tau_p &= (1 - \delta_p) \varphi_2 \sigma_p + \delta_p \varphi_1 \sigma_p \end{aligned} \tag{7}$$

and

$$\begin{aligned} \varphi_2 \sigma_{p+1} &= (1 - \theta_p) \varphi_2 \sigma_p + \theta_p \varphi_1 \tau_p \\ \varphi_2 \tau_p &= \delta_p \varphi_2 \sigma_p + (1 - \delta_p) \varphi_1 \sigma_p \end{aligned} \tag{8}$$

where $\sigma_0 = 20$, $\tau_0 = 30$, $\theta_0 = 0.70$ and $\delta_0 = 0.80$.

Here in the following table, we compare four different cases of this iterative method:

Table 3.1: Comparison of iteration of $\varphi_2 \sigma_{p+1}$

Steps	Iteration (5)	Iteration (6)	Iteration (7)	Iteration (8)
1	20	20	20	20
2	4.648383871	3.543260003	3.543260004	4.648383188
3	3.032758079	3.182386197	3.182386198	3.032758080
4	1.982147798	2.686616150	2.686616151	1.980700986
5	1.460445409	2.251688906	2.251688907	1.460311366
6	1.216632658	1.909824677	1.909824679	1.216592446
7	1.103196627	1.653251704	1.653251631	1.103184565
8	1.049808611	1.465354602	1.465354552	1.049804993
9	1.024304786	1.329760594	1.329760560	1.024303702
10	1.011956989	1.232793443	1.232831915	1.011956664
11	1.005915998	1.163953511	1.163980442	1.005915902
12	1.002938283	1.115266094	1.115284946	1.002938255
13	1.001462984	1.080935480	1.080948677	1.001462977
14	1.000729583	1.056779417	1.056788655	1.000729583
15	1.000364205	1.039807876	1.039814343	1.000364205
16	1.001892300	1.027896654	1.027932324	1.000181923

Emphasis of Coefficients on the Convergence Rate of Fixed Point Iterative Algorithm in Banach Space

17	1.000090907	1.022366281	1.019583767	1.000090907
18	1.000045437	1.015664181	1.013724207	1.000045437
19	1.000033789	1.010968819	1.009614730	1.000022714
20	1.000014678	1.007680119	1.006734203	1.000011550
21	1.000006674	1.005377057	1.004715888	1.000005677
22	1.000003137	1.003764426	1.003302095	1.000002838

Table 3.2: Comparison of iteration of $\varphi_2 \tau_n$

Steps	Iteration (5)	Iteration (6)	Iteration (7)	Iteration (8)
1	4.120592286	3.065961283	4.120592300	3.065961285
2	2.621470658	3.257556507	2.400446023	4.141655056
3	1.769923962	2.836752044	1.799849601	2.849154969
4	1.361156399	2.390474630	1.502050077	1.875403875
5	1.165478769	2.020978859	1.328848724	1.409430803
6	1.081676801	1.737443965	1.220301834	1.192901692
7	1.029590004	1.527337400	1.149594475	1.092131876
8	1.019381602	1.374637844	1.102487564	1.044580031
9	1.009557113	1.264982086	1.070646665	1.021797125
10	1.004736039	1.186820715	1.048910226	1.010738969
11	1.002354662	1.131515575	1.035322900	1.005318682
12	1.001173174	1.092359204	1.024319399	1.002643370
13	1.000585302	1.064821535	1.016820691	1.001316708
14	1.000292256	1.045460106	1.011673147	1.000656816
15	1.000146007	1.031704586	1.008120561	1.000327936
16	1.000072967	1.025552812	1.005665307	1.000163823

17	1.000036472	1.017897595	1.003956174	1.000081868
18	1.000021002	1.012533631	1.002764551	1.000040921
19	1.000011331	1.008776198	1.001932801	1.000020457
20	1.000005222	1.006144667	1.001351768	1.000010227
21	1.000002478	1.004301931	1.000945641	1.000005113

From the above tables, we analyze the iteration (5) converges more rapid than all iterations (6)-(8). The following pie graph represents the convergence behaviour of this iterative method:




Fig. 3.1: Graphical representation of convergence speed

III. CONCLUSION

The above analysis of convergence speed of particular iteration process, it is investigated that the convergence rate of such iterative processes vary efficiently when the coefficients involved in these methods are interchanged with its respective iterative schemes. By numerical consideration, clearly the iterative process (5) converges with a rate faster than the iterations (6)–(8). Moreover, it is clear from the graphical presentation that the iterative procedure (7) moves quicker than the iteration (6) and the iteration (8) approaches to its fixed point better than the iterations (6) and (7). So all in all, we say that the iterative plan (5) moves more rapid than all the cases of fixed point iterative process considered in this manuscript. Therefore, it is analyzed that the coefficients of these iterative processes have a notable effect on the convergence rate on these schemes. Following is the convergence behaviour of this iterative procedure:




Fig. 2: Convergence Behaviour Iteration Scheme

IV. REFERENCES

- [1] Ishikawa S., Fixed points by a new iteration method. Proceedings of the American Mathematical Society, 44, 147–150(1974)
- [2] Ishikawa S., Fixed points and iteration of a nonexpansive mapping in a Banach space. Proceedings of the American Mathematical Society, 59(1), 65–71(1976)
- [3] Talman L.A., Fixed points for considering multi functions in metric spaces with convex structure. Kodai Math, Sem. Rep. 29, 62-70 (1977).
- [4] Osilike M. O., Stability results for Ishikawa fixed point iteration procedure. Indian J. Pure Appl. Math., 26, No. 10, 937-341 (1995)
- [5] Xu B., Noor M. A., Fixed-Point Iterations for Asymptotically Non expansive Mappings in Banach Spaces. Journal of Mathematical Analysis and Applications, 267, 444–453 (2002)
- [6] Berinde V., Picard iteration converges faster than the Mann iteration for a class of quasi-contractive operators. Fixed Point Theory Applications, (2), 97-105(2004)
- [7] Kima T. H., XuH.K.,Stron, Convergence of Modified Mann iterations. Nonlinear Analysis 61, 51– 60 (2005)
- [8] Babu G.V.R., and Vara Prasad K.N.V.V., Mann iteration converges faster than Ishikawa iteration for the class of Zamfirescu operators. Fixed Point Theory Applications, Article ID 49615, 1-6 (2006)
- [9] Popescu O., Picard iteration converges faster than Mann iteration of quasi-contractive operators. Mathematical Communications 12(2), 195-202 (2007)
- [10] ManikumariN. and Murugappan A., Fuzzy Logic Based Model for Optimization of Tank Irrigation System, Journal of Engineering and Applied Sciences, Medwell Journals, 3 (2), 199-202, (2008)
- [11] Alfred O. B., Strong Convergence results for the Jungck–Mann iteration (2010).

- [12] Akewe H. Olaoluwa H., On the Convergence of Modified three-step Iteration process for generalized Contractive-like operators. *Bulletin of Mathematical Analysis and Applications*, Vol. 4 Issue 3, 78-86 (2012)
- [13] ZhiqunXue and Guiwen Lv., The convergence of the modified Mann and Ishikawa iterations in Banach spaces. *Journal of Inequalities and Applications*, Article Number: 188 (2013)
- [14] Chen J. and Wu D., Convergence theorems of modified Mann Iterations. *Fixed Point Theory and Applications*, 282-285 (2013)
- [15] Juan Xiao, Lei Deng, and Ming-ge Yang, Convergence of Modified Multi-Step Iterative for a finite family of Asymptotically Quasi-nonexpansive mappings. *Commun. Korean Math. Soc.* 29, No. 1, 83–95 (2014).
- [16] Fathollahi S., Ghiura A., Postolache M. and Rezapour S., A comparative study on the convergence rate of some iteration methods involving contractive mappings. *Fixed Point Theory Appl.* (2015)
- [17] Renu Chug and Rekha Rani, A Weak Convergence Theorem for Variational Inequalities and Fixed Point Problems in a Real Hilbert Space. *Journal of Engineering and Applied Sciences*, Medwell Journals, 2962-2970 (2016)
- [18] ThakurB. S., Thakur D., and PostolacheM., A New Iteration Scheme for Approximating Fixed Points of Nonexpansive Mappings. *Filomat*, 30(10), 2711–2720, (2016)
- [19] Chauhan S.S., Utreja K., Imdad M. and Ahmadullah Md., Strong Convergence Theorems for a Quasi-contractive type Mapping Employing a new Iterative Scheme with an Application. *Honam Mathematical J.* 39, No. 1, 1-25(2017)
- [20] Kumar N., and Chauhan Gonder S.S., Analysis of Jungck-Mann and Jungck-Ishikawa Iteration schemes for their speed of convergence. *AIP Publishing*, Vol. 2050 (2018)
- [21] Kumar N., and Chauhan Gonder S.S., A Review on the convergence speed in the Agarwal et al. and Modified-Agarwal Iterative Schemes. *Universal Review*, Vol. 7, Issue X, 163-167(2018)
- [22] Kumar N., and Chauhan Gonder S.S., An Illustrative Analysis of Modified-Agarwal and Jungck-Mann Iterative Procedures for their Speed of Convergence. *Universal Review*, Vol. 7, Issue X, 168-173 (2018)
- [23] Kumar N., and Chauhan Gonder S.S., Examination of the Speed of Convergence of the Modified-Agarwal Iterative Scheme. *Universal Review*, Vol. 7, Issue X, 174-179 (2018)
- [24] Zena Hussein Maibed, Some Generalized n-Tuplet Coincidence Point Theorems for Non-Linear Contraction Mappings, *Journal of Engineering and Applied Sciences*, Medwell Journals, 13 (24): 10375-10379, (2018)
- [25] Kumar N., and Chauhan Gonder S.S., Speed of convergence examined by exchange of coefficients involved in Modified-Ishikawa iterative scheme. *Future Aspects in Engineering Sciences and Technology*, Vol. 2, Chandigarh University, 440-447 (2018)

- [26] Kumar N., and Chauhan Gonder S.S., Self-Comparison of Convergence Speed in Agarwal, O'Regan and Sahu's S-Iteration. International Journal on Emerging Technologies, 10 (2b), 105-108 (2019)